### FINAL REPORT

### NASA Grant NAG5-2754

NOVEMBER 1, 1996 - OCTOBER 31, 1999

### DATA REDUCTION AND ANALYSIS FROM THE SOHO SPACECRAFT

F. M. Ipavich, Project Director

Submitted by the

Department of Physics University of Maryland College Park, MD 20742

Joe Gurman, Project Scientist NASA, GSFC

### Instrumentation

The Mass-determining Time-of-Flight (MTOF) instrument is a modified, improved version of the "MASS" instrument flown on the WIND mission, and was built by the University of Maryland (UMD). The unique entrance system was built at the University of Bern (UBE) in Switzerland. Along with the Charge Determining Time-of-Flight (CTOF) and Suprathermal Time-of-Flight (STOF) instruments, MTOF is a part of the Charge, Element, and Isotope Analysis System (CELIAS). The Mass-determining Time-of-Flight sensor on SoHO is designed for the precise determination of the chemical and isotopic composition of the solar wind over a wide range of solar wind conditions. MTOF can measure the chemical and isotopic composition of the solar wind for most elements with mass  $\leq 70$  or so. MTOF uses electrostatic analysis followed by a time-of-flight measurement in a specially designed harmonic potential.

The Proton Monitor (PM) is a subsensor of the MTOF instrument, which is one of the 3 time-of-flight instruments comprising the CELIAS experiment. MTOF determines high-resolution mass spectra of heavy solar wind ions and uses a very wide bandwidth energy-per-charge analyzer to maximize counting statistics. The PM was designed to assist in the interpretation of MTOF data and for that reason uses a similar wide bandwidth analyzer that limits the accuracy of derived solar wind parameters.

#### Health of Instrument

We routinely monitor the health of the MTOF instrument, and support the various spacecraft maneuvers with appropriate command sessions to reduce our high voltages during thruster burns, followed by re-establishing the nominal mode for MTOF.

Every day at noon, a pair of programs is automatically run on the latest files received from the Experiment Operations Facility (EOF). These programs write housekeeping and rate information to ASCII files that are then sent via electronic mail to several members of our group. Visual examination of these files provides a check on instrument health, and could give rapid warning if an instrument problem were to arise. Another program that is automatically run at noon each day generates a binary file of rate information suitable for analysis on a Macintosh.

In order to monitor the health of the instrument and the quality of the data in the longer term, six programs are run each day that read the most recent Level Zero file. Two of these programs generate files of rate information for both the MTOF main sensor and the PM subsensor. Two others generate files containing instrument status and housekeeping information. Another program finds the minimum and maximum values of various housekeeping quantities, and still another generates a binary file of rate information suitable for analysis on a Macintosh. The ASCII files generated by the first five programs are stored on line and also printed and kept in binders. This allows easy access to housekeeping, status, and rate information for the entire mission.

Flight operations for the CELIAS experiment during the reporting period (November 1996 to October 1999) have included the following:

Safing and recovery of the MTOF, STOF, PM, and CTOF sensors during SoHO operations which involve the use of thrusters (station keeping, momentum management, and reaction wheel maintenance) These operations take place approximately every 3 months, and typically require commanding over a 4-5 day period.

Reactivation of the MTOF, STOF, PM, and CTOF sensors after the loss and recovery of the SoHO spacecraft in 1998.

Turning MTOF back on after instrument "latchups" caused the sensor to be turned off. This has happened three times to date.

Special commanding has been required for CTOF (which began experiencing problems in August 1996), and for software patches to the DPU (patches 12, 13, 14, and 15). The purpose of patch #15 is to deal with the aforementioned instrument "latchups". During a latchup condition, the main MTOF current rises to abnormally high levels. Since a latchup could occur while SoHO is not being tracked, it is possible that such high currents could persist for hours, if the patch were not in place. This would probably cause severe damage to the sensor. Patch 15 allows the CELIAS DPU to monitor the MTOF main current. If this current exceeds a given threshold value four times consecutively (indicating a latchup condition) the DPU will turn the MTOF sensor off for 5 minutes. This should clear the latchup. Then the DPU will turn the sensor back on in STANDBY mode (all high voltages off).

New TSTOL procedures have been created for the DPU, MTOF and STOF.

Documentation on the CELIAS ESR response and the MTOF emergency procedures has been supplied to the Science Operations Coordinator (SOC).

CELIAS housekeeping is processed daily and sent to MPAe, MPE, and UMD. We have responded to the FOT regarding yellow and red flags observed for CTOF, STOF, and MTOF parameters.

## Solar Wind Web Page

We have also spent considerable effort in establishing and maintaining a page on the World Wide Web that presents almost real-time solar wind data from our MTOF/PM subsensor on the SoHO spacecraft. It was announced in an on-line newsletter (SPA section newsletter, Vol. 3, Issue 66) that is circulated to the space physics community. Presently, the PM page receives about 700 "hits" per day. The URL is:

http://umtof.umd.edu/pm

The PM web page has been continuously revised and improved. The algorithms used to derive solar wind parameters from the PM data were refined and tested over a time period of months, and the organizational structure of the web page was substantially changed. In addition, the solar wind flow angle out of the ecliptic plane was added to the list of parameters that are plotted.

The web page contains plots of solar wind parameters for the most recent 2 days of data and the most recent 2 weeks of data. The solar wind parameters plotted are bulk speed, proton density, thermal speed, and angle out of the ecliptic plane. The current location of the SoHO spacecraft, taken from predicted orbit files, is also displayed on the page.

Links on the page lead to: descriptions of the plotted parameters and the PM sensor; older data organized by Carrington Rotation; older orbit data; an "interesting figures" page where we have expanded scale plots of e.g. shocks, high speed streams, low density time periods, etc; and an incomplete "shock table" listing solar, interplanetary, and magnetospheric conditions near the time of shocks observed by the PM. The older PM data is available in the form of tables of ASCII data and also in the form of plots. The older orbit data is a plain text table.

Data files are transferred electronically via FTP from the EOF to the UMTOF system at Maryland. The PM page is updated automatically within a few

minutes of the receipt of a file from the EOF. The data is processed using a FORTRAN program and new plots are generated by IDL as needed; human intervention is not required.

SoHO is never in the magnetosphere and enjoys excellent data coverage. The data on the Web page are typically between a few minutes and a few hours old.

We have also responded to numerous requests for PM data made by members of the scientific community around the world.

## **Data Analysis**

A large number of programs have been written to assist in data analysis. These programs are described in Appendix C. To facilitate use of these programs, considerable effort was made to provide online documentation available via the World Wide Web. This documentation consists of almost 300 files, containing over 2.3 megabytes of information.

An automated routine (called "Shockspotter") that searches the PM data for shocks is in development, and is currently being tested. The algorithm is being fine-tuned to minimize the number of false positives and false negatives.

### Science

We continue to analyze and interpret data from the MTOF sensor on SoHO, using recently obtained calibration data (from the MTOF spare) to improve our understanding of instrument response. We have presented a number of talks demonstrating the excellent instrument resolution and also displaying the temporal behavior of solar wind Fe<sup>54</sup> and Fe<sup>56</sup> isotopes and the element Cr(52). In addition, the isotopic ratios Ne<sup>20</sup>/Ne<sup>22</sup> = (13.8  $\pm$  0.7) and Ne<sup>20</sup>/Ne<sup>21</sup> = (440  $\pm$ 110) were obtained from MTOF data. These ratios agree with the values obtained from the Apollo foil solar wind experiments and with values that have been derived from measurements on lunar and meteoritic samples.

Peter Bochsler (CELIAS P.I.) of the University of Bern in Switzerland spent his sabbatical at the University of Maryland. Professor Bochsler assisted in the development of MTOF analysis procedures, lending his expertise in several areas. These included charge state yields and scattering of ions traversing the Carbon foil in the TOF sensor, and the theoretically expected abundances and charge states of ions in different solar wind flow types.

In order to derive more precise instrument efficiencies, we took the MTOF spare sensor to the University of Bern in Switzerland for additional calibration. There were two trips. The first was undertaken in 1998. The duration of the trip was from March 21 to April 3 1998. The CASYMS accelerator at Bern is capable of producing beams of a variety of elements and energies. We used beams of He<sup>+</sup>, O<sup>+</sup>, and Ar<sup>+</sup> with beam energies ranging from as low as 0.7 keV for Helium to as high as 60 keV for Argon. By removing the deflection system for a portion of the calibration, we were able to obtain information about the contributions to the overall efficiency due to the time-of-flight system alone, as well as the sensor as a whole. The second trip lasted from October 1 to October 21 1999. The MEFISTO accelerator at Bern is capable of producing beams of a variety of elements, energies, and charge states, including such elements as iron and calcium, which are unavailable at other accelerators. By removing the deflection system for the calibration, we obtained information about the contributions to the overall efficiency due to the time-of-flight system alone.

Talks on CELIAS were given at a variety of venues, including the AAAS Meeting in Seattle, IAGA in Uppsala, the Solar Wind 9 Meeting in Nantucket, the AGU Spring Meetings in Baltimore and Boston, the 7th and 8th SoHO Workshops (in Northeast Harbor, Maine and in Paris, respectively), and the 24th EGS General Assembly in The Hague. Papers have been submitted to the Journal of Geophysical Research, to Solar Physics, and to GRL. The talks and papers are listed in a complete bibliography accompanying this report as Appendix D.

The first determination of the Na abundance in the solar wind was reported at the Spring AGU meeting. Na with its 5.12 V First Ionization Potential (FIP) is the most extreme low-FIP element observed in the solar wind to date. The proximity of Na to Ne in mass as well as the similar and narrow charge state distributions of Na and Ne in the corona render these two elements as almost ideal markers to accurately identify small variations between low- and high-FIP solar wind with CELIAS/MTOF. The short first ionization time of Na and the relation of the solar wind Na abundance to a very well known photospheric value make it a prime candidate for investigating coronal-hole associated FIP fractionation and for sensitive testing of FIP/FIT models. The behavior of Na relative to low-FIP and high-FIP elements was compared, for both interstream and coronal-hole associated solar wind flows. Preliminary results indicated that Na behaves as expected for a low-FIP element in these two types of solar wind flows.

The first determination of the Al abundance in the solar wind, obtained with CELIAS/MTOF, was also reported at the AGU meeting. The analysis covered two short periods in November 1998. The solar wind Al abundance is particularly interesting, because of the extremely short ionization time of this element under normal chromospheric conditions. The Al abundance has been evaluated in comparison with the neighboring low-FIP element Mg. A preliminary evaluation yielded an Al/Mg abundance ratio of 0.088 for the interstream period and a ratio of 0.086 for a period of coronal hole associated solar wind. Within the estimated uncertainties of less than 20% these ratios are consistent with each other, and also compatible with the photospheric and the meteoritic abundance ratios. At the same time the normal bias between low- and high-FIP elements is confirmed for the periods of investigation. Hence, Al, which ionizes much faster than Mg under chromospheric conditions, shows no deviation in its abundance from the other low-FIP elements for the two investigated periods.

In addition, we made a pair of collaborative studies, together with the SWICS sensor on the ACE spacecraft. In the first of these, we used combined measurements from ACE/SWICS and SOHO/MTOF to investigate the onset of a well defined coronal hole-associated high speed stream on DOY 69, 1998. The SWICS O7/O6 ratio shows that the transition from slow to high-speed solar wind flow was relatively abrupt (less than 2 hours), while the transition back to slow wind was much broader (a few days). The density pile-up region preceding the coronal hole flow had typical low-speed charge states. The precipitous drop in the Oxygen ionization temperature was accompanied by an increase in the solar wind bulk and thermal speeds, a decrease in density, and an enhanced velocity and temperature difference between the solar wind Protons and Alpha Particles. The Fe isotope ratios remained unchanged throughout these transitions. In the second collaboration, we used combined measurements from ACE/SWICS and SOHO/MTOF to investigate the behavior of Iron during the unusual CME of May 2 and 3, 1998. We found a surprisingly large distribution of Fe charge states, ranging from +3 to at least +16, with the lowest and highest charge states occasionally observed simultaneously. To our knowledge these are the first measurements of Fe+3 in the solar wind. In contrast to the charge state distributions, the Fe isotope ratio remained constant with a value very close to its meteoritic ratio.

# Appendix A: Routine SOHO Tasks

This document lists those automated SOHO science tasks that are performed on a regular basis on the UMTOF system at the University of Maryland.

The science oriented tasks, which consist mainly of the procedure SOHO\_DAILY being run in one its two modes (SUITE1 and SUITE2), are performed daily.

Another set of tasks is done regularly and with varying frequency to maintain the Proton Monitor (PM) home page on UMTOF. For details on how the output of these procedures and programs are put together, see a separate article on how the PM home page is constructed.

# Routine SOHO Science Tasks

Below is a list of tasks performed on a regular basis. Unless noted otherwise the task is performed once each day.

Task	What Is Done
SOHO_DAILY (LATESTLZ)	Run a suite of programs to process the latest SOHO LZ files and
SOHO_DAILY (LATESTEOF)	create binary files which can be viewed on a Mac. Run a suite of programs to process the latest SOHO REL/QKL file and create a Mac binary file for it. Mail ASCII output of
RECENT48	programs to selected users. Perform other housekeeping tasks. Check if there are any new quicklook data. If so, generate a new 48-hour PM plot and update the PM home page. This checks for new files every five minutes.
RECENT_2WEEK	Generate 14-day PM plot, update PM home page.
CRN	Generate a new or updated plot of PM data for a Carrington
LOG_SOHO_POSITION_NEW	Rotation period. Update web pages to point to new plot. This runs approximately every 14 days.  Obtain and record the SOHO spacecraft's position information and other information. These items of data are used by
SPLIT_QKL	RECENT48 and RECENT_2WEEK.  Locate data gaps in QKL (quicklook) data files, split QKL files
31 LIT_QKL	into smaller files without gaps. This is done every 12 hours.
MASTER_CH	Change format of recent (within the last 3 days) Command History (CH) files, extract certain Command History records, and append them to the file MASTER_CH.DAT
CHECK_ROLL_NEW	Read latest attitude file and check if roll angle is outside acceptable limits.
SOHO_NIGHTLY	Perform various file maintenance tasks, such as purging and deleting certain files.

# Appendix B: SOHO Science Procedures on the UMTOF System

This document gives information on SOHO data processing procedures used at the University of Maryland at College Park UMTOF system. It is divided into three sections:

# Detail Description of SOHO Procedures

Procedure: SOHO\_DAILY (LATESTLZ)

Purpose:

Process recent SOHO LZ files, creating ASCII and Mac binary files. Write all output files to

DISKM4:[DAILY]. Print all ASCII output files to UMTOF printer.

When run: each day at 06:00

Programs run:

EDB process latest LZ science file, create Mac binary file
DUMMY process latest LZ science file, create ASCII output file
RATES process latest LZ science file, create ASCII output file
PMRATES process latest LZ science file, create ASCII output file
HK process latest LZ HK file, create ASCII output file

HK MINMAX process latest LZ HK file, create ASCII output file

Data sets:

Science and housekeeping LZ files copied to the system since 6:00 of the preceding day. Specifically, all .DAT1 files (both G029 and G013) in SOHO\_LZ\_DATA: whose modification date is later than 6:00 of the preceding day.

Procedure: SOHO\_DAILY (LATESTEOF)

Purpose:

Process latest EOF data, create Mac binary file, extract rate and housekeeping information, and email it to selected personnel.

When run:

each day at 12:00

Programs run:

EDB process latest EOF science, create Mac binary file RATES process latest EOF science, create ASCII output file PK process latest EOF science, create ASCII output file

Data sets:

Latest science EOF file (CELSC1\_\*.\*) located in SOHO\_EOF\_DATA: with size at least 2500 blocks, and latest housekeeping (CELHK\_\*.\*) EOF file located in SOHO\_EOF\_DATA with size at least 50 blocks. The latest data set is the last one reported in a DIRECTORY listing.

Procedure: CHECK\_ROLL\_NEW

Purpose:

Examine the attitude files and detect if the spacecraft's roll angle has exceeded its allowable range.

When run: each day at 23:30

Programs run:

CHECK\_ROLL

Data sets:

SOHO definitive attitude files (SOHO\_AT\_DATA:SO\_AT\_DEF\*.CDF) copied to the system since 23:30 of the preceding day. DISKM1:[SOHO]ROLL.DAT contains log of when satellite roll angle was outside of norm.

Appendix B: SOHO Science Procedures on the UMTOF System

Procedure: LOG\_SOHO\_POSITION\_NEW

Purpose:

Examine the orbit files, extract the spacecraft's position, and append it to a file. Also extract and write out the distance from the sun, the space latitude and longitude and the current Carrington Rotation number (earth). Finally, create an HTML file of the orbit information.

When run: each day at 23:30

Programs run:

LOG\_SOHO\_POSITION

Data sets:

Predicted orbit files (SOHO\_OR\_DATA:SO\_OR\_PRE\*.CDF) copied to UMTOF since 23:30 of the preceding day. The modification date is used to select the data sets. Previously this procedure used definitive orbit files, but they are no longer generated.

Description:

The quantities GSE position (X, Y, and Z) in Earth radii, Space latitude and longitude in degrees, and Carrington Rotation Number (earth) are extracted from the second entry of each orbit file and appended to the file [SOHO]ORBIT.DAT. In addition the distance from the sun is computed and written to the file.

The file OLDORBIT.HEADER, the file ORBIT.DAT mentioned above, and the file OLDORBIT.TRAILER and concatenated and copied to the file OLDORBIT.HTML in the PM WWW area.

Procedure: MASTER\_CH

Purpose:

Change the file characteristics of recently received Command History files so they can be easily examined. Extract all command history records dealing with CELIAS and append them to a master CELIAS command history log.

When run: each day at 00:00

Programs run:

SET FILE SEARCH

**APPEND** 

Data sets:

All command history files (SOHO\_CH\_DATA:SO\_CH\_NUL\*.DAT) whose format is not StreamLF. The file format is changed to StreamLF. The file is then searched for records containing the strings " CEL" or " FB". Those records are extracted from the command history file and appended to the file SOHO\_CH\_DATA:MASTER\_CH.DAT .

Procedure: RECENT48

Purpose:

This procedure runs as a batch job every 5 minutes. If there is new QKL or REL data it runs the program PMSW to extract the most recent 48 hours of data. It then invokes the procedure PMPLOT.COM and passes it the filename and several other parameters, requesting that it update the PM home page.

When run: every  $\hat{5}$  minutes

Programs run:

PMSW IDL

**GET\_PREDICT\_POSITION** 

PMPLOT.COM

TIMES.PRO

PMPLOT\_TICKS.PRO

PMPLOT\_IO.PRO

PMPLOT\_GRAPH.PRO

PMPLOT\_ADJUST.PRO

### Appendix B: SOHO Science Procedures on the UMTOF System

PMPLOT\_RANGES.PRO PMPLOT\_INIT.PRO PMPLOT\_MAIN.PRO

Data sets:

Realtime and quicklook Celias science files from the EOF facility

(SOHO\_EOF\_DATA:CELSC1\*.\*) which are larger than 10 blocks and represent data from the 48 hour period just ended. Slightly more or less data may be displayed depending on the phase of the starting time of the interval (always starts on multiple of 4 hours) and the presence (or rather the absence) of data sets near then end of the interval.

Procedure: RECENT\_2WEEK

Purpose:

This procedure runs as a batch job every morning at 7:00. It runs the program PMSW to extract the most recent 14 days of data. It then invokes the procedure PMPLOT.COM and passes it the filename and several other parameters, requesting that it update the PM home page.

When run: each day at 07:00

Programs run:

PMSW IDL

PMPLOT.COM

TIMES.PRO

PMPLOT\_TICKS.PRO

PMPLOT\_IO.PRO

PMPLOT\_GRAPH.PRO

PMPLOT\_ADJUST.PRO

PMPLOT RANGES.PRO

PMPLOT\_INIT.PRO

PMPLOT MAIN.PRO

Data sets:

Realtime and quicklook Celias science files from the EOF facility and Level Zero (LZ) science files from the ISTP facility which represent data from the 14 day interval just ended. Intervals always start at 0000 GMT.

Procedure: CRN

Purpose:

Generate plots of PM quantities based on Carrington Rotation instead of an arbitrary calendar period. Each plot is generated four times as additional/more reliable data becomes available.

When run:

approximately every 14 days at 1:23

more precisely, it is run once after the start of a Carrington Rotation period and once at the midpoint of the rotation period

Programs run:

CRN\_PLOT.COM

**PMSW** 

IDL.

PMPLOT.COM

TIMES.PRO

PMPLOT\_TICKS.PRO

PMPLOT\_IO.PRO

PMPLOT\_GRAPH.PRO

PMPLOT\_ADJUST.PRO

PMPLOT RANGES.PRO

PMPLOT\_INIT.PRO

PMPLOT\_MAIN.PRO

Data sets:

Each time CRN is run two plots are generated. In each case the interval (i.e., rotation period) of the first plot generated exactly matches the interval of a plot generated earlier. It replaces the earlier plot in the CRN web page.

Description:

In mode A the first plot (period AX, file suffix \_A3) uses mostly Level Zero science files, but may use some realtime and quicklook Celias science files for the rotation period preceding the current period. The second plot (period AY, file suffix \_A1) uses only realtime and quicklook Celias science files for the current rotation period. This generates a plot of a partial period.

In mode B the first plot (period BX, no file suffix) uses only Level Zero science files for the rotation period preceding the one just ended. The second plot (period BY, file suffix \_B2) uses a combination of Level Zero, realtime, and quicklook science Celias files science files for the rotation period just ended.

Procedure: SOHO\_NIGHTLY

Purpose:

Perform various maintenance tasks, such as deleting old files or data sets, and purging files which have multiple versions. Also perform certain monthly maintenance tasks, such as deleting old LZ data sets and resetting the version numbers of certain frequently used files.

When run:

each day at 3:33

Programs run:

**PURGE** 

DELETE

Data sets:

Nightly:

LZ, OR, AT, CH files which have multiple VMS versions - purge

All \*.SFDU files - delete

All CELSC1 EOF files that are smaller than 10 blocks but not currently being transferred - delete All EOF files that are more than 45 days old - delete

LZ, OR, AT files which have multiple data set versions - delete earlier version numbers

All 2-day graphs more than 2 days old - delete

All 2-week graphs more than 14 days old - delete

All full 2-week graphs and their associated image map files more than 1 day old - delete, but keep composite graphs

Monthly:

RECENT48 log files, its associated control files, PMSW.DAT, and

PMSW.USED - purge and reset version number

Procedure: SPLIT\_QKL

Purpose:

Scan the EOF directory for Quicklook (\*.QKL) files. For each file examine it for large data gaps (more than I hour). Create another, potentially two, output files to eliminate the gap. Two output files are created if there is a significant (more than 5 minutes) amount of data preceding the gap. Rename all processed files into another directory for archival purposes. Change file attributes on created files so PMSW will be able to process them.

When run: every 4 hours

Programs run:

SPLIT\_QKL

Data sets:

SOHO EOF DATA:\*.OKL

Creates corresponding output files with user-specified file type; e.g.., \*.MOD

# Appendix C: SOHO Programs on UMTOF

This appendix describes most of the SOHO programs in use at the University of Maryland on the UMTOF system. The following notes apply to all programs:

- All executables reside in DISKM1:[SOHO.EXE], which is pointed to by the logical name SOHO\_EXE: .
- Working source code for the programs is archived in DISKM1:[SOHO.SOURCE], which is pointed to by the logical name SOHO\_SOURCE: .
- Source code for the object libraries which are used by the programs is archived in DISKM1:[SOHO.LIB.SOURCE].
- All output files generated by these programs which have a file type of .DAT are suitable for loading into a spreadsheet program such as KaleidaGraph. In addition they have the time in one of the columns.
- All output files generated by these programs which have a file type of .TABLE are more free format and can not be used in a spreadsheet.
- Both .DAT and .TABLE output files contain ASCII information.
- Output files having a file type of .BIN are binary files with stream-of-bytes format suitable for reading on a Mac.
- The following programs allow spanning multiple files: DUMMY, EDB, EDBPM, HEAT\_CURR, HK, HK\_FULL, PHATREND, PMRATES, PMRSTACK, PMRSTACKA, PMRT, PMSW, RATES, THERM2, and THERM4
- The programs that span files handle nested files appropriately.
- Most of the programs can be forced to use a specific input file by defining a logical name. This is especially useful in the event of a spacecraft problem. If the logical name SOHO\_INPUT\_FILE is defined, the programs will not prompt for a start or stop time, instead processing the single file specified by the logical name. The programs for which this does *not* work are HOUR\_AVERAGE, LOG\_SOHO\_POSITION, PMOFFLINE, PMSIMEOF, PMSW\_REFORMAT, SAMPLE\_SW\_SMS, SPLIT\_QKL, STACK, and SWE\_AVERAGE.

# Summary of SOHO Science Programs On UMTOF

Below is a list of the programs and a brief description of each. This is followed by a more detailed description of each program.

Program Name	Purpose
C_SPEED	computes CTOF speed
DUMMY	dumps dummy EDBs
DUMP_PACKET	dumps packets
DUMP_TIMES	dumps times
EDB	reads and optionally dumps EDBs
EDBPM	reads and dumps EDBs with solar wind parameters added
HEAT_CURR	dumps currents and heater status
HK	dumps some quantities from HK packets
HK_FULL	dumps HK packets (all quantities)
HK_MINMAX	finds lowest and highest values of certain MTOF-related quantities
HOUR_AVERAGE	averages 5-minute resolution files, producing 1-hr and 2-hr
	averages
LOG_SOHO_POSITION	records position of SOHO satellite
PHATREND	counts PHA words in user selected region
PMOFFLINE.COM	generates PM plots without changing PM home page
PMRATES	dumps PM rates and PM E/Q values
PMRSTACK	generates stacked plots of PM counts vs. radius

### Appendix C: SOHO Programs on the UMTOF System

PMRSTACKA generates stacked plots of PM counts vs. radius, with altered radius

bins

PMRT dump distributions in radius and theta of PM counts
PMSIMEOF Given 6 PM rates, write SW parameters to screen
fits PM data and derives solar wind parameters

PMSW DEFORMAT

PMSW\_REFORMAT Reformats data files produced by PMSW

**RATES** 

SAMPLE\_SW\_SMS samples WIND SMS solar wind data files

SPLIT\_QKL detects large gaps in QKL files and splits them into one or two files

without gaps

dumps rates

STACK DCL command file to run either PMRSTACK or PMRSTACKA

SWE\_AVERAGE reads SWE KP files, generates 1-hr and 6-hr averages

THERM2 dumps temperatures from SVM HK2 THERM4 dumps temperatures from SVM HK4

# Detailed Description of SOHO Science Programs on UMTOF

C\_SPEED This program opens a CELIAS science data file from disk or CD, reads

and dumps the file header, and then reads science data packets, forms EDB's and then SR's. It then prints the CTOF speed, the time, and CTOF\_STATUS\_LEN provided that CTOF speed is not 25.0 km/s. This

program does not span files.

DUMMY

This program opens a CELIAS science data file from disk or CD, reads

and dumps the file header, and then reads science data packets, looking for dummy EDB's. Data from Dummy EDB's is stored in a huge array. Portions of that data (MTOF classification tables, levels and other mode-specific information, voltage steps, etc.) are interpreted, then displayed. The output file DUMMY.TABLE contains roughly 1 page per 8 hour

period.

DUMP\_PACKET This program dumps the time and the source sequence counter for each

packet. If a sync word is present that is dumped as well. The output file PACKET.DAT contains one line per packet. This program does not span

files.

DUMP\_TIMES This program opens a CELIAS science data file from disk or CD, reads

and dumps the file header, and then reads science data packets, forms EDB's and then SR's. It then dumps the packet times for each EDB 0, the SR start time from D\_Spare, and the difference between them. The output file TIMES.DAT contains a few lines per science record. This program

does not span files.

EDB This program opens a CELIAS science data file from disk or CD, reads

and dumps the file header, and then reads science data packets, forms EDB's and then SR's. It can dump each science record to EDB.TABLE, producing 1 page per 15 seconds, or it can dump the information to a binary file suitable for use on a Mac, or it can do both. If the input file is an LZ file, the binary file is named YY\_DOY\_MMDD.BIN. If the input

file is an REL/QKL file the binary file is named

YY\_DOY\_MMDDHH.BIN. In both cases, the year, day of year, month, and day specified by YY, DOY, MM, and DD come from the input file

name, as does the hour, specified by HH in the latter case.

EDBPM This program opens a CELIAS science data file from disk or CD, reads

and dumps the file header, and then reads science data packets, forms

EDB's and then SR's. It dumps the information to a binary file, together with solar wind parameters calculated from the data. If the input file is an LZ file, the binary file is named YY\_DOY\_MMDD.BIN. If the input file is an REL/QKL file the binary file is named YY\_DOY\_MMDDHHP.BIN. In both cases, the year, day of year, month,

and day specified by YY, DOY, MM, and DD come from the input file name, as does the hour, specified by HH in the latter case.

HEAT\_CURR

This program dumps currents and heater status from SVM HK1. It can deal with faulty times (caused, e.g., by a satellite reset) by checking if the year of the first packet is before the launch date (1995). If so the starting Epoch time is set to zero (early 1958), and the ending Epoch time is left at its current value (usually the current date). As a result, files are still spanned, but every file header is printed - not just the header of the first data file, as usual. The output file is HEAT\_CURR.DAT.

HK

This program opens a CELIAS housekeeping file from disk or CD, reads and dumps the file header, and then reads and dumps HK packets. It prints only a few quantities (voltages, currents, temperatures, and a few others), and prints them in a column format to facilitate the use of KaleidaGraph. The output file is named HK.DAT, and can only be printed using the MP4 command (its output is 148 columns across!).

HK\_FULL

This program opens a CELIAS housekeeping file from disk or CD, reads and dumps the file header, and then reads and dumps HK packets. The output file, named HK.TABLE, will contain one page per 15 seconds.

HK\_MINMAX

This routine finds the two lowest values, the two highest values, and the average for DPU and MTOF-related quantities from the Celias Housekeeping files. It operates on a single LZ file and does not span files. Its output is written to the file HK\_MINMAX.DAT.

HOUR AVERAGE

This program reads the data files produced by PMSW at 5 minute resolution and generates 1 hour and 2 hour resolution files. These files have the same name as the input file, but with the extents .1HR and .2HR, respectively. The version of HOUR\_AVERAGE in [SOHO.EXE] reads files with 8 columns of data, exclusive of the date and time. The version in [SOHO.TEST] reads files with 12 columns of data exclusive of the date and time.

LOG\_SOHO\_POSITION

This program reads a predicted or definitive SOHO orbit CDF file and extracts several quantities at just after midnight. It then appends these to the file ORBIT.DAT in the directory [SOHO]. The quantities extracted from the orbit file are: GSE position (X, Y, and Z), printed in Earth radii; Space latitude and longitude, printed in degrees; and Carrington Rotation Number (earth). The program, when run, prompts the user for an orbit (OR) filename. These files are located in a directory pointed to by the logical names SOHO\_OR\_DATA and OR.

**PHATREND** 

This program allows the user to select up to 10 TOF regions and an interval of summation. It then counts the number of PHA words with TOF's in each region, and outputs the result along with the input file header to a file named PHATREND.DAT. If the regions overlap then an event may be counted more than once. If a file named PHATREND.INPUT exists in the default directory then inputs will be used from it.

read from it.

Appendix C: SOHO Programs on the UMTOF System

PMRATES This program dumps all six PM rates and PM E/Q values in a column

format readable by KaleidaGraph. The output file is named

PMRATES.DAT.

PMRSTACK This program generates stacked plots of the PM counts vs. radius on either

logarithmic or linear scales. The output file is named PMRSTACK.DAT.

PMRSTACKA This program generates stacked plots of the PM counts vs. radius on either

logarithmic or linear scales. A different binning scheme than the original

is used. The output file is named PMRSTACK.DAT.

PMRT This program dumps the distributions in radius and theta of PM counts.

The output file is named PMRT.TABLE.

PMSIMEOF This program (formerly named TABLE\_PM\_E) prompts the user for six

PM rates. It then calculates the solar wind speed, density, and thermal

speed using the same algorithm as PMSWA.

PMSW This program fits PM data and derives solar wind speed, density, thermal

speed, and N/S angle. The output file is named PMSW.DAT. The most

recent version (in [SOHO.EXE]) produces 23 columns of output.

PMSW\_REFORMAT This program reads a file produced by PMSW and reformats it, printing

out solar wind speed, density, thermal speed, and flow angle, along with orbital data from the CDF files. Definitive orbit data are used when available; Predictive orbit data are used when necessary. The output file may be either HTML or plain text, as the user desires. The user chooses

both the output file's name and its format.

RATES This program opens a CELIAS science data file from disk or CD, reads

and dumps the file header, and then reads science data packets, forms EDB's and then SR's. It then dumps only the rates, printing them in a column format to facilitate their use in KaleidaGraph. The output file is named RATES.DAT and can only be printed using the command MP4 (its

output is 168 columns across!).

SAMPLE\_SW\_SMS This program samples all files of the form SW\_96xxx.DAT (where xxx is

a day of year) in the directory SMS1:[COLLIER.SW\_SPEED]. Every 20th proton speed is printed out (along with its corresponding time) to the

file SMS\_SPEED.DAT.

SPLIT\_QKL This program scans the EOF directory for SOHO Quicklook (\*.QKL)

files. For each QKL file the program reads the file and determines if it contains a large time gap. If so the QKL file is split into two smaller QKL files: the first has all the data up to the start of the gap, and the second has all data after the gap. The output files are appropriately named to reflect their start times and are given a file type of .MOD. If the amount of data in the first output would be less than a certain threshold, typically five minutes, it is not created and only the second file is created. If there is no gap or the gap is fairly short a copy of the entire file is made; the new file is given a file type of .MOD. In either case, after the copy operation(s) is

complete, the input file is moved to another directory.

Both the minimum length (in time) of the first data segment and the size of

the time gap are parameters which can be easily changed. They are

currently set at 5 minutes and 30 minutes, respectively.

### Appendix C: SOHO Programs on the UMTOF System

STACK

Not technically a program, STACK is a DCL command file that prompts the user for input and then invokes PMRSTACK, invokes PMRSTACKA, or uses a preexisting file generated by an earlier invocation of these programs to produce a stacked plot.

SWE\_AVERAGE

This program reads the SWE KP files at 90 second resolution and generates 1 hour and 6 hour resolution files. These files have the same name as the input file, but with the extents .1HR and .6HR, respectively.

THERM2

This program dumps temperatures from SVM HK2. It uses a polynomial fit. This program can deal with faulty times (caused, e.g., by a satellite reset). When s/c times erroneously indicate a year before 1995, files are still spanned, but every file header is printed - not just the header of the first data file, as usual. The output file is THERM2.DAT.

THERM4

This program dumps temperatures from SVM HK4. It uses a polynomial fit. This program can deal with faulty times (caused, e.g., by a satellite reset). When s/c times erroneously indicate a year before 1995, files are still spanned, but every file header is printed - not just the header of the first data file, as usual. The output file is THERM4.DAT.

# Appendix D: List of Publications Relevant to the CELIAS Instrument

#### Table of Contents

Talks and Poster Presentations Papers on Instrument Topics Papers on Scientific Topics Related Presentations

### Talks and Poster Presentations

Model for the Mass Fractionation in the January 6, 1997, CME Wurz, P., Bochsler, P., and Lee, M. A., 1999 Fall AGU in California, USA.

Determination of the argon isotopic ratio of the solar wind using SOHO/CELIAS/MTOF J.M. Weygand, F.M. Ipavich, P. Wurz, J. A. Paquette, and P. Bochsler, 8th SOHO Workshop in Paris, France.

Magnesium Isotopic Abundance in the Slow and Coronal Hole Associated Solar Wind: SOHO/CELIAS/MTOF Measurements. H. Kucharek, F.M. Ipavich, R. Kallenbach, B. Klecker, H. Grünwaldt, M. R. Aellig, and P. Bochsler, 8th SOHO Workshop in Paris, France.

Elemental and ionic composition of suprathermal ions in the May 2-3, 1998 CME K. Bamert, R.F. Wimmer-Schweingruber, R. Kallenbach, M. Hilchenbach, B. Klecker, A. Bogdanov, 1999 Spring AGU in Boston, U.S.A.

Magnesium isotopic abundance in slow and coronal hole associated solar wind: SOHO/CELIAS/MTOF measurements, H. Kucharek, F. M. Ipavich, R. Kallenbach, B. Klecker, H. Gruenwaldt, M. R. Aellig, P. Bochsler, Spring AGU 1999 in Boston, USA

Determination of the argon isotopic ratio of the solar wind using SOHO/CELIAS/MTOF J.M. Weygand, F.M. Ipavich, P. Wurz, J.A.Paquette, and P. Bochsler, 1999 Spring AGU in Boston, U.S.A.

Determination of the argon isotopic ratio of the solar wind using SOHO/CELIAS/MTOF J.M. Weygand, F.M. Ipavich, P. Wurz, J.A.Paquette, and P. Bochsler, EGS Conference in Den Haag Netherlands, 1999.

Elemental and Ionic Composition of Suprathermal Ions in the May 2-3, 1998 CME K. Bamert, R.F. Wimmer-Schweingruber, R. Kallenbach, M. Hilchenbach, B. Klecker, A. Bogdanov, EGS Conference in Den Haag, Netherlands, 1999.

Isotopic Composition of Magnesium Measured with SOHO/CELIAS/MTOF In Slow and

Coronal Hole Associated Solar Wind, H. Kucharek, F. M. Ipavich, R. Kallenbach, B. Klecker, H. Grünwaldt, M. R. Aellig, and P. Bochsler, the 24th General Assembly in Den Hague, Netherlands, April 19-23, 1999.

Nonlinear Analysis of Solar Wind Data from the Proton Monitor Onboard SOHO, R. Sachs, H. Kucharek, F. M. Ipavich, R. Kallenbach, B. Klecker, and P. Bochsler, the 24th General Assembly in Den Hague, Netherlands, April 19-23, 1999.

The Si/O Abundance Ratio in the Solar Wind Measured with CELIAS/CTOF and the Search for the FIP Effect in Coronal Hole Associated Wind. M.R. Aellig, H. Holweger, S. Hefti, P. Wurz, H. Grünwaldt, F.M. Ipavich, and P. Bochsler, the AGU fall meeting 1998.

The Iron, Silicon, and Oxygen Abundance in the Solar Wind Measured with SOHO/CELIAS/MTOF, P. Wurz, M.R. Aellig, F.M. Ipavich, S. Hefti, P. Bochsler, and A.B. Galvin, the AGU fall meeting 1998.

Isotopic Composition of Solar Wind Nitrogen: First In-Situ Determination by CELIAS/MTOF Onboard SOHO R. Kallenbach, J. Geiss, F. M. Ipavich, G. Gloeckler, P. Bochsler, F. Gliem, S. Hefti, M. Hilchenbach, and D. Hovestadt, the AGU fall meeting 1998.

The Fe/O Elemental Abundance Ratio in the Solar Wind, M.R. Aellig, P. Bochsler, P. Wurz, H. Grünwaldt, S. Hefti, H. Holweger, F.M. Ipavich, and B. Klecker, the Solar Wind 9 Conference in Nantucket, MA, USA, 5-9 October 1998.

CELIAS/CTOF Measurement of O<sup>5+</sup>: A Key to the Maze of Coronal Electron Velocity Distribution, H. Grünwaldt, M. Hilchenbach, E. Marsch, P. Bochsler, and F. Ipavich Solar Wind 9 Conference in Nantucket, MA, USA, 5-9 October 1998.

Highly unusual behavior of Solar Wind heavy ions in early May 1998, F. M. Ipavich, G. Gloeckler, J. Paquette, T. Zurbuchen, P. Bochsler, P. Wurz, Solar Wind 9 Conference in Nantucket, MA, USA, 5-9 October 1998.

First Measurements of Sulfur Isotopes in the Solar Wind, A.B. Galvin, F.M. Ipavich, J. Paquette, R. Kallenbach, B. Klecker, and P. Wurz, Solar Wind 9 in Nantucket, MA, USA. 5-9 October 1998.

SOHO/CELIAS Measurements of Sulfur Isotopes in the Solar Wind, A.B. Galvin, F.M. Ipavich, J. Paquette, D. Hovestadt, P. Wurz, and R. Kallenbach, 7th SOHO Workshop, 28 September - 1 October 1998, Northeast Harbor, Maine, USA.

Temporal Behavior of Solar Wind Fe and O in High Speed Stream Onsets, F. M. Ipavich, G. Gloeckler, T. Zurbuchen, S. Hefti, P. Bochsler, L. A. Fisk, R. Kallenbach, J. Paquette, and R. F. Wimmer-Schweingruber, 7th SOHO Workshop, 28 September - 1 October 1998, Northeast Harbor, Maine, USA.

Heliospheric Energetic Neutral Hydrogen Atoms as Observed with SOHO/CELIAS, M. Hilchenbach, K.C. Hsieh, D. Hovestadt, B. Klecker, H. Grünwaldt, F.M. Ipavich, P. Bochsler, and the CELIAS Team, at the COSPAR meeting, July 12-19, 1998, Nagoya, Japan.

SOHO Solar Extreme Ultraviolet Full Disk Flux Observations Since SOHO SEM Commissioning in December 1995, D. L. Judge, 1st TIGER Symposium, Freiburg Germany, June 18/19, 1998 Measurements of the solar EUV/UV radiation

Multispacecraft Analysis of a <sup>3</sup>He Rich Solar Energetic Particle Event with ACE/SEPICA and SOHO/HSTOF, A.T. Bogdanov, E. Möbius, B. Klecker, M. Hilchenbach, D. Hovestadt, L.M. Kistler, M.A. Popecki, E.J. Lund, D. Heirtzler, A.B. Galvin, P. Bochsler, H. Grünwaldt, F.M. Ipavich, and F. Gliem, AGU Spring Meeting, 26-29 May 1998, Boston, MA, USA.

Fractionation of Si, Ne, and Mg Isotopes in the Solar Wind as Measured by SOHO/CELIAS/MTOF, R. Kallenbach, F.M. Ipavich, H. Kucharek, P. Bochsler, J. Geiss, F. Gliem, G. Gloeckler, H. Grünwaldt, M. Hilchenbach, and D. Hovestadt, AGU Spring Meeting, 26-29 May 1998, Boston, MA, USA.

Iron Isotopes in the Solar Wind: Measurements From SoHO/CELIAS/MTOF, F.M. Ipavich, J.A. Paquette, P. Bochsler, R. Kallenbach, P.Wurz, H. Grünwaldt, M. Hilchenbach, and D. Hovestadt, AGU Spring Meeting, 26-29 May 1998, Boston, MA, USA.

Elemental Composition of the January 6, 1997, CME, P. Wurz, P. Bochsler, M.R. Aellig, F.M. Ipavich, A.B. Galvin, AGU Spring Meeting, 26-29 May 1998, Boston, MA, USA.

The Fe/O Elemental Abundance in the Solar Wind as Determined From SOHO/CELIAS/CTOF Measurements, M.R. Aellig, S. Hefti, H. Grünwaldt, P. Bochsler, P. Wurz, F.M. Ipavich, M. Hilchenbach, D. Hovestadt, AGU Spring Meeting, 26-29 May 1998, Boston, MA, USA.

The Influence of Suprathermal Electrons on the Derivation of Coronal Electron Temperatures from Solar Wind Minor Ion Charge State Spectra Obtained from SOHO/CELIAS/CTOF, M.R. Aellig, P. Bochsler, H. Grünwaldt, S. Hefti, P. Wurz, M. Hilchenbach, D. Hovestadt, F.M. Ipavich, F. Gliem, at the EGS 23rd General Assembly 20-24 April, 1998, Nice, France.

Solar Wind Isotopic Abundance Ratios of Ne, Mg, and Si Measured by SOHO/CELIAS/MTOF as a Diagnostic Tool for the Inner Solar Corona, R. Kallenbach, F.M. Ipavich, H. Kucharek, P. Bochsler, J. Geiss, F. Gliem, G. Gloeckler, A.B. Galvin, H. Grünwaldt, M. Hilchenbach, D. Hovestadt, at the EGS 23rd General Assembly 20-24 April, 1998, Nice, France.

The Search with CELIAS/CTOF for O<sup>5+</sup> as a Solar Wind Constituent, Heiner Grünwaldt, Martin Hilchenbach, Eckhardt Marsch, Peter Bochsler, and the CELIAS Team, at the EGS 23rd General Assembly 20-24 April, 1998, Nice, France.

Correlated C+ and O+ Source Near 0.2 AU, Heiner Grünwaldt, Ingrid Mann, Martin Hilchenbach, Peter Bochsler, and the CELIAS Team, at the EGS 23rd General Assembly 20-24 April, 1998, Nice, France.

The Silicon, Oxygen, and Iron Abundance in the Solar Wind, P. Wurz, F.M. Ipavich, P. Bochsler, M.R. Aellig, S. Hefti, R. Kallenbach, A.B. Galvin, H. Grünwaldt, M. Hilchenbach, F. Gliem, D. Hovestadt, and the CELIAS Team, at the EGS 23rd General Assembly 20-24 April, 1998, Nice, France.

Absolute solar EUV irradiances from the CELIAS/SEM during the long-term minimum of solar cycle 23, H. S. Ogawa, D. L. Judge, D. R. McMullin, and P. Gangopadhyay AGU Fall Meeting, December 12, 1997, San Francisco, CA

Solar Wind Abundance of Magnesium Isotopes From SOHO/CELIAS/MTOF, H. Kucharek, R. Kallenbach, P. Bochsler, F. M. Ipavich, M. Hilchenbach, H. Grünwaldt, F. Gliem, B. Klecker, D. Hovestadt and the CELIAS team, AGU Fall Meeting, 8-12 December 1997, San Francisco, CA, USA.

Solar Wind Abundance of Magnesium Isotopes from SOHO/CELIAS/MTOF, H. Kucharek, R. Kallenbach, P. Bochsler, F. M. Ipavich, M. Hilchenbach, H. Grünwaldt, F. Gliem, B. Klecker, D. Hovestadt, and the CELIAS team, at AGU Fall Meeting, 8-12 December 1997, San Francisco, CA, USA.

Pickup Ions in the Mass Interval 20-30 AMU Observed with SOHO/CELIAS/CTOF vs. Model Calculations of Distribution Functions Accounting for Heliospheric and Interstellar Seed Populations, A.T. Bogdanov, B. Klecker, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, E. Möbius, F. M. Ipavich, F. Gliem, P. Bochsler, at AGU Fall Meeting, 8-12 December 1997, San Francisco, CA, USA.

Charge State Composition of Energetic Helium as Observed in CIRs at 1 AU with SOHO/CELIAS, M. Hilchenbach, H. Grünwaldt, H. Kucharek, B. Klecker, D. Hovestadt, R. Kallenbach, P. Bochsler, F. Gliem, A. Galvin, K. Chotoo, F.M. Ipavich and the CELIAS team, at AGU Fall Meeting, 8-12 December 1997, San Francisco, CA, USA.

Solar Wind Cross Correlation Measurements between SOHO And WIND, N. Silverman, M. Coplan, F. Ipavich, A. Galvin, K. Ogilvie, A. Lazarus, J. Steinberg, AGU Fall Meeting, 8-12 December 1997, San Francisco, CA, USA.

Solar Wind Minor Ion Charge States Observed With High Time Resolution With CELIAS/CTOF, M.R. Aellig, H. Grünwaldt, P. Bochsler, P. Wurz, S. Hefti, R. Kallenbach, F.M. Ipavich, D. Hovestadt, M. Hilchenbach, and the CELIAS Team, 31st ESLAB Symposium, 22-25 September 1997, ESTEC, Noordwijk, The Netherlands.

Limits on the Fractionation of Isotopes in the Solar Wind as Observed with SOHO/CELIAS/MTOF, R. Kallenbach, P. Bochsler, F.M. Ipavich, A.B. Galvin, R. Bodmer, S. Hefti, H. Kucharek, F. Gliem, H. Grünwaldt, M. Hilchenbach, B. Klecker, D. Hovestadt, and the CELIAS Team, 31st ESLAB Symposium, 22-25 September 1997, ESTEC, Noordwijk, The Netherlands.

Energetic Neutral Hydrogen Atoms Distribution in the Heliosphere: Observations with SOHO/CELIAS, M. Hilchenbach, K.C. Hsieh, H. Grünwaldt, F.M. Ipavich, F. Gliem, P. Bochsler, B. Klecker, D. Hovestadt, and the CELIAS TEAM, 31st ESLAB Symposium, 22-25 September 1997, ESTEC, Noordwijk, The Netherlands.

The Silicon, Oxygen, and Iron Abundance in the Solar Wind, P. Wurz, P. Bochsler, M.R. Aellig, S. Hefti, R. Kallenbach, F.M. Ipavich, A.B. Galvin, H. Grünwaldt, M. Hilchenbach, F. Gliem, D. Hovestadt, and the CELIAS Team, 31st ESLAB Symposium, 22-25 September 1997, ESTEC, Noordwijk, The Netherlands.

Magnesium Isotope Composition In The Solar Wind as Observed with SOHO/CELIAS/MTOF, H. Kucharek, F.M. Ipavich, R. Kallenbach, F. Gliem, H.

Grünwaldt, M. Hilchenbach, B. Klecker, P. Bochsler, D. Hovestadt, and the CELIAS Team, 5th SOHO Workshop, 17-20 June, 1997, Oslo, Norway.

Solar Wind Iron Charge States Observed with High Time Resolution with CELIAS/CTOF, M.R. Aellig, H. Grünwaldt, P. Bochsler, P. Wurz, S. Hefti, R. Kallenbach, F.M. Ipavich, D. Hovestadt, M. Hilchenbach, and the CELIAS Team, 5th SOHO Workshop, 17-20 June, 1997, Oslo, Norway.

Highlights from CELIAS, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F.M. Ipavich, M.R. Aellig, W.I. Axford, H. Balsiger, A. Bogdanov, A. Bürgi, M.A. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, S. Hefti, K.C. Hsieh, D.L. Judge, R. Kallenbach, B. Klecker, H. Kucharek, M.A. Lee, Y. Litvinenko, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, H.S. Ogawa, K. Reiche, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, 5th SOHO Workshop, 17-20 June, 1997, Oslo, Norway.

Oxygen Freeze-in Temperature Measured by SOHO/CELIAS, S. Hefti, P. Bochsler, H. Grünwaldt, M. Aellig, D. Hovestadt, M. Hilchenbach, F. M. Ipavich, R. Kallenbach, P. Wurz, and the CELIAS team, AGU Spring Meeting, 27-30, May, 1997, Baltimore, USA.

How well is the Solar Isotopic Composition Reflected in the Solar Wind?, P. Bochsler, R. Bodmer, R. Kallenbach, P. Wurz, F.M. Ipavich, A.B. Galvin, G. Gloeckler, M. Hilchenbach, H. Grünwaldt, D. Hovestadt, B. Klecker, AGU Spring Meeting, 27-30, May, 1997, Baltimore, USA.

Solar Wind Measurements from SoHO during the CME events of early 1997, F.M. Ipavich, A.B. Galvin, G. Gloeckler, S. Lasley, J. Paquette, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F. Gliem, and the CELIAS Team, IAGA 97, August 4-15, 1997, Uppsala, Sweden.

Distribution of Energetic Neutral Hydrogen Atoms as Observed with SOHO/CELIAS, M. Hilchenbach, K.C. Hsieh, H. Grünwaldt, F.M. Ipavich, F. Gliem, P. Bochsler, B. Klecker, D. Hovestadt, and the CELIAS TEAM, IAGA 97, August 4-15, 1997, Uppsala, Sweden.

Solar Wind Measurements from SoHO During the CME Events of Early 1997, F.M. Ipavich, A.B. Galvin, P. Bochsler, H. Grünwaldt, M. Hilchenbach, D. Hovestadt, F. Gliem, and the CELIAS Team, AGU Spring Meeting, 27-30, May, 1997, Baltimore, USA.

Results on Neon Pickup Ions From Measurements With SOHO/CELIAS/CTOF And Comparison With Model Calculations, A. T. Bogdanov, B. Klecker, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F. M. Ipavich, F. Gliem, P. Bochsler, E. Möbius, and the SOHO/CELIAS TEAM, EGS General Assembly, 1997, Vienna, Austria.

New SoHO Observations: The Solar Wind in Situ, A.B. Galvin, F.M. Ipavich, H. Grünwaldt, M. Hilchenbach, F. Gliem, P. Bochsler, D. Hovestadt, and the CELIAS Team, AAAS meeting, February 16, 1997, Seattle, WA, USA.

Observation of Energetic Neutral Hydrogen Atoms with SOHO/CELIAS, M. Hilchenbach, K.C. Hsieh, H. Grünwaldt, F.M. Ipavich, F. Gliem, P. Bochsler, D. Hovestadt, and the CELIAS Team, EGS General Assembly, 1997, Vienna, Austria.

Charge Composition in Solar Energetic Particle Events as Observed with SOHO/CELIAS at 1 AU, B. Klecker, A. Bogdanov, H. Kucharek, D. Hovestadt, M. Hilchenbach, H. Grünwaldt, R. Kallenbach, P. Bochsler, F. Gliem, F.M. Ipavich, and the CELIAS team, EGS General Assembly, 1997, Vienna, Austria.

Helium Charge Composition in CIRs as observed with SOHO/CELIAS at 1 AU, M. Hilchenbach, H. Grünwaldt, R. Kallenbach, P. Bochsler, H. Kucharek, D. Hovestadt, F. Gliem, F.M. Ipavich, and the CELIAS team, EGS General Assembly, 1997, Vienna, Austria.

CELIAS-CTOF on SOHO as a Fast Remote Coronal Thermometer, H. Grünwaldt, P. Bochsler, D. Hovestadt, M. Hilchenbach, F.M. Ipavich, and F. Gliem, EGS General Assembly, 1997, Vienna, Austria.

Crossing the Tail of Venus with CELIAS-CTOF on SOHO, H. Grünwaldt, M. Neugebauer, P. Bochsler, D. Hovestadt, M. Hilchenbach, F.M. Ipavich, and F. Gliem, EGS General Assembly, 1997, Vienna, Austria.

The Silicon and Oxygen Abundance in the Solar Wind Measured with CELIAS/MTOF, P. Wurz, R. Kallenbach, P. Bochsler, M.A. Aellig, S. Hefti, F.M. Ipavich, A.B. Galvin, G. Gloeckler, H. Grünwaldt, M. Hilchenbach, F. Gliem, D. Hovestadt, and the CELIAS collaboration, EGS General Assembly, Vienna, Austria (April, 1997).

Magnesium Isotope Abundance Measured with SOHO/CELIAS/MTOF, H. Kucharek, F.M. Ipavich, R. Kallenbach, F. Gliem, H. Grünwaldt, M. Hilchenbach, B. Klecker, P. Bochsler, D. Hovestadt, and the CELIAS Team, EGS General Assembly, Vienna, Austria (April, 1997).

First year solar EUV irradiance from SOHO by the CELIAS/SEM, H. S. Ogawa, D. L. Judge, D. R. McMullin, P. Gangopadhyay, and A. B. Galvin, EGS XXII GENERAL ASSEMBLY Vienna, 21-25 APRIL 1997 Session ST5: Poster

Solarer Wind und energiereiche Teilchen: Erste Ergebnisse des CELIAS Experimetes auf SOHO, B. Klecker, H. Grünwaldt, M. Hilchenbach, F.M. Ipavich, F. Gliem, P. Bochsler und D. Hovestadt, annual meeting of the Deutsche Physikalische Gesellschaft, München, Germany, March (1997).

Ein schnelles Fernthermometer für die Corona: Messungen mit CELIAS-CTOF auf SOHO, H. Grünwaldt, D. Hovestadt, P. Bochsler, M. Hilchenbach, F. Ipavich and the CELIAS Team, annual meeting of the Deutsche Physikalische Gesellschaft, Munich, Germany, März (1997).

SOHO/CELIAS-CTOF: Ein Flug durch den Schweif der Venus, H. Grünwaldt, M. Neugebauer, P. Bochsler, M. Hilchenbach, F. Ipavich, D. Hovestadt, and the CELIAS collaboration, annual meeting of the Deutsche Physikalische Gesellschaft, Munich, Germany, März (1997).

Measurements and Modeling of Ne Pickup Ions Based on Data from the SOHO/CELIAS/CTOF Instrument, A.T. Bogdanov, H. Grünwaldt, M. Hilchenbach, F.M. Ipavich, F. Gliem, P. Bochsler, D. Hovestadt, B. Klecker, E. Möbius, Kollaboration-SOHO/CELIAS TEAM, annual Meeting of the Deutsche Physikalische Gesellschaft, München, Germany, März (1997).

Messung der Magnesium Isotopenhäufigkeit im solaren Wind mit SOHO/CELIAS/MTOF, H. Kucharek, F.M. Ipavich, F. Gliem, H. Grünwaldt, M. Hilchenbach, B. Klecker, P. Bochsler, D. Hovestadt und die CELIAS-Kollaboration, annual Meeting of the Deutsche Physikalische Gesellschaft, München, Germany, März (1997).

Beobachtung von energetischem neutralem Wasserstoff und Helium mit dem Experiment CELIAS auf der SOHO Mission, M. Hilchenbach, J. Hsieh, D. Hovestadt, P. Bochsler, H. Grünwaldt, F.M. Ipavich und die SOHO/CELIAS Kollaboration, annual Meeting of the Deutsche Physikalische Gesellschaft, München, Germany, März (1997).

Detection of Energetic Neutral Hydrogen Atoms of Heliospheric Origin, M. Hilchenbach, K.C. Hsieh, P. Bochsler, D. Hovestadt, H. Grünwaldt, F.M. Ipavich, W.I. Axford, H. Balsiger, A. Bürgi, M.A. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, D.L. Judge, R. Kallenbach, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, H.S. Ogawa, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, AGU Fall Meeting, San Francisco, CA, USA, (1996).

Radiometric Results Obtained to Date with the SOHO Extreme Ultraviolet Monitor (SEM) and a Sounding Rocket Underflight, D.L. Judge, Joint PTB/ESA Seminar on 'Spectroradiometry with the Solar and Heliospheric Observatory (SOHO) First Results and Future Work', Berlin Germany, October 31-Nov. 1, 1996 Solar EUV Monitor (SEM) experiment on SOHO (to review current calibration results)

First Solar EUV Irradiances from SOHO by SEM, Judge, D.L., D.R. McMullin, H.S. Ogawa, D. Hovestadt, M. Hilchenbach, E. Moebius, L.R. Canfield, R. Watts, M. Khne, and P. Wurz, 17th International Workshop/SOLERS22 1996 Workshop, June 17-21, 1996 National Solar Observatory/Sacramento Peak, Sunspot, New Mexico, USA Group 4&5: contributed

Solar Corona Diagnostic with Solar Wind Iron Charge Spectra, M.R. Aellig, H. Grünwaldt, S. Hefti, P. Wurz, P. Bochsler, W.I. Axford, H. Balsiger, A. Bürgi, M.A. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, M. Hilchenbach, D. Hovestadt, K.C. Hsieh, F.M. Ipavich, D.L. Judge, R. Kallenbach, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, H.S. Ogawa, K.U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, Fall Meeting of the Swiss Physical Society, Zürich, Switzerland, (1996).

Solar Wind Observations with SOHO/CELIAS, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F.M. Ipavich, K.-U. Reiche, W.I. Axford, H. Balsiger, A. Bürgi, M. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, M. Hilchenbach, K.C. Hsieh, D.L. Judge, R. Kallenbach, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, H.S. Ogawa, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, EPS Fall Meeting, Sevillia, Spain (1996).

Solar Wind Composition: First Results from SOHO and Future Expectations, A. B. Galvin, F. M. Ipavich, G. Gloeckler, M. Coplan, D. Hovestadt, M. Hilchenbach, A. Bürgi, B. Klecker, M. Scholer, P. Bochsler, H. Balsiger, J. Geiss, R. Kallenbach, P. Wurz, H. Grünwaldt, W. I. Axford, S. Livi, E. Marsch, B. Wilken, F. Gliem, K.-U. Reiche, M. A. Lee, E. Möbius, K. C. Hsieh, M. Neugebauer, G. G. Managadze, M. I. Verigin, 188th Meeting of the American Astronomical Society, Madison, Wisconsin, USA, (1996).

CELIAS on SOHO: Solar Wind and Suprathermal Ion Measurements at L1, D. Hovestadt, M. Hilchenbach, A. Bürgi, B. Klecker, P. Laeverenz, M. Scholer, H. Grünwaldt, W.I.

Axford, S. Livi, E. Marsch, B. Wilken, P. Winterhoff, P. Bedini, M.A. Coplan, A.B. Galvin, G. Gloeckler, P. Bochsler, H. Balsiger, J. Fischer, J. Geiss, R. Kallenbach, P. Wurz, K.-U. Reiche, F. Gliem, D.L. Judge, K.H. Hsieh, E. Möbius, M.A. Lee, G.G. Managadze, M.I. Verigin, M. Neugebauer, Western Pacific Geophysics Meeting, Brisbane, Australia, (July 23-27,1996).

CELIAS on SOHO - Solar Particle Abundances at L1, D. Hovestadt, P. Bochsler, H. Grünwaldt, M. Hilchenbach, F.M. Ipavich, K.-U. Reiche, W.I. Axford, H. Balsiger, A. Bürgi, M. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, K.C. Hsieh, R. Kallenbach, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, M. Scholer, M.I. Verigin, B. Wilken, P. Wurz, 31st COSPAR Scientific Assembly, Birmingham, United Kingdom, (14-21 July, 1996).

Solar Particle Composition on SOHO: First Measurements from the CELIAS Experiment, F.M. Ipavich, D. Hovestadt, P. Bochsler, H. Grünwaldt, M. Hilchenbach, K.-U. Reiche, W.I. Axford, H. Balsiger, A. Bürgi, M. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, K.C. Hsieh, R. Kallenbach, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, AGU Spring Meeting, Baltimore, MD, USA, EOS (1996) \$205.

Initial results of the Solar EUV Monitor (SEM) Irradiance Measurements Aboard SOHO, D.L. Judge, D.R. McMullin, H.S. Ogawa, D. Hovestadt, M. Hilchenbach, E. Möbius, L.R. Canfield, R. Watts, M. Kühne, and P. Wurz, AGU Spring Meeting, Baltimore, MD, USA, EOS (1996) S214.

Isotopic Composition of Solar Wind Neon Derived from Observations with CELIAS/MTOF, R. Kallenbach, F.M. Ipavich, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, K.-U. Reiche, W.I. Axford, H. Balsiger, A. Bürgi, M. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, M. Hilchenbach, K.C. Hsieh, D.L. Judge, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, H.S. Ogawa, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, AGU Spring Meeting, Baltimore, MD, USA, EOS (1996) S215.

## Papers on Instrument Topics

CELIAS: The Charge, Element, and Isotope Analysis System for SOHO, D. Hovestadt, M. Hilchenbach, A. Bürgi, B. Klecker, P. Laeverenz, M. Scholer, H. Grünwaldt, W.I. Axford, S. Livi, E. Marsch, B. Wilken, P. Winterhoff, F.M. Ipavich, P. Bedini, M.A. Coplan, A.B. Galvin, G. Gloeckler, P. Bochsler, H. Balsiger, J. Fischer, J. Geiss, R. Kallenbach, P. Wurz, K.-U. Reiche, F. Gliem, D.L. Judge, K.H. Hsieh, E. Möbius, M.A. Lee, G.G. Managadze, M.I. Verigin, M. Neugebauer, *Solar Physics*, 162, 441, 1995.

The Charge, Element, and Isotope Analysis System CELIAS on SOHO, D. Hovestadt, P. Bochsler, H. Grünwaldt, F. Gliem, M. Hilchenbach, F.M. Ipavich, D.L. Judge, W.I. Axford, H. Balsiger, A. Bürgi, M. Coplan, A.B. Galvin, J. Geiss, G. Gloeckler, K.C. Hsieh, R. Kallenbach, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, Lecture Notes in Physics, edts. A.O. Benz and A. Krüger, *Proc. of the CESRA* 

workshop held in Caputh/Potsdam, Germany, 16-20 May 1994, 271-278, Springer-Verlag, Berlin and Heidelberg, Germany, 1995.

Charge exchange of B, C, O, Al, Si, S, F and Cl passing through thin carbon foils at low energies: Formation of negative ions, R. Kallenbach, M. Gonin, P. Bochsler, and A. Bürgi, *Nucl. Instr. and Meth. B* 103, 111, 1995.

Charge Exchange of Low Energy Particles Passing Through Thin Carbon Foils: Dependence on Foil Thickness and Charge Yields of Mg, Ca, Ti, Co and Ni, M. Gonin, R. Kallenbach, P. Bochsler, and A. Bürgi, *Nucl. Instr. and Meth. B101*, 313, 1995

Reflection of Ultraviolet Light at 121.6nm From Rough Surfaces, T. Zurbuchen, P. Bochsler, and F. Scholze, *Opt. Eng. 34*, 1303, 1995.

Charge Exchange of Atoms with High First Ionization Potentials in Carbon Foils at E<35keV/u, M. Gonin, R. Kallenbach, and P. Bochsler, *Nucl. Instr. & Meth. B* 94, 15, 1994.

Charge Exchange of Hydrogen Atoms in Carbon Foils at 0.4-120 keV, M. Gonin, R. Kallenbach, and P. Bochsler, *Rev. Sci. Instr.* 65, 648, 1994.

Calibrations of Entrance Systems of Two Mass Spectrometers for Solar Wind Measurements, S. Hefti and R. Kallenbach, *Proc. of the Week of Doctoral Students*, Charles University, Prague, Czech Republic 176, 1994.

Charge Exchange of Hydrogen Ions in Carbon Foils, R. Kallenbach, M. Gonin, A. Bürgi, and P. Bochsler, *Nucl. Instr. & Meth. B* 83, 68, 1994.

Charge Exchange of Low-Energy Ions in Thin Carbon Foils II. Results for Ions of B, C, F, Ne, Na, Si, S, Cl, Ar, K, and Fe, A. Bürgi, M. Gonin, M. Oetliker, P. Bochsler, J. Geiss, T. Lamy, A. Brenac, H.J. Andrä, P. Roncin, H. Laurent, and M.A. Coplan, J. Appl. Phys. 73, 4130, 1993.

Interaction of Solar Wind Ions With Thin Carbon Foils - Calibration of Time-Of-Fight Spectrometers, M. Gonin, A. Bürgi, M. Oetliker, and P. Bochsler, *Proc. of the first SOHO workshop on coronal streamers, coronal loops, and coronal and solar wind composition*, Annapolis, Maryland, USA, ESA SP-348, 381, 1992.

Copper Sulfide Coating to Reduce Ion and UV-Light Specular Reflections in Spacecraft-Borne Mass Spectrometers, W. Saxer, M. Wüst, T. Zurbuchen, and P. Bochsler, *Proc. Surfair IX*, Cannes, edition Ampère, 1992.

Simulation of Ion Backscattering from Rough Surfaces, M. Wüst and P. Bochsler, *Nucl. Instrum. Methods in Phys. Res. B* 71, 314, 1992.

Partial Discharge Measurements to Determine Long-Term Characteristics of Thin Epoxy Plates under Cyclic Stress, H.J. Weber, K. Lehmann, P. Bochsler, and J. Fischer, ESA SP-320, 335, 1991.

Ion Composition Measurements, A. Bürgi, Adv. Space Res. 11(1), 391, 1991.

Charge Exchange of Low-Energy Ions in Thin Carbon Foils, A. Bürgi, M. Oetliker, P. Bochsler, J. Geiss, M.A. Coplan, *J. Appl. Phys.* 68, 2547, 1990.

The CELIAS Experiment Onboard the SOHO Spacecraft, P. Bochsler, *Proc. of the IVth International Seminar: Manufacturing of Scientific Space Instrumentation*, V.M. Balebanov (ed.), Academy of Sciences, Space Research Institute, Frunze, 137, 1989.

CELIAS - Charge, Element, and Isotope Analysis System for SOHO, D. Hovestadt, J. Geiss, G. Gloeckler, E. Möbius, P. Bochsler, F. Gliem, F.M. Ipavich, B. WIlken, W.I. Axford, H. Balsiger, A. Bürgi, M. Coplan, H. Dinse, A.B. Galvin, K.I. Gringauz, H. Grünwaldt, K.C. Hsieh, B. Klecker, M.A. Lee, G.G. Managadze, E. Marsch, M. Neugebauer, W. Rieck, M. Scholer, and W. Stüdemann, ESA SP-1104 69, 1989.

## Papers on Scientific Topics

Determination of the Abundance of Aluminum in the Solar Wind with SOHO/CELIAS/MTOF, P. Bochsler, F. M. Ipavich, J. A. Paquette, J. M. Weygand, and P. Wurz, *J. Geophys. Res.* (in press).

Abundances and Charge States of Particles in the Solar Wind, P. Bochsler, submitted to *Reviews of Geophysics*.

Enhanced Solar Wind 3He<sup>2+</sup> Associated with Coronal Mass Ejections. Ho G.C., Hamilton D.C, Gloeckler G., and Bochsler P., *Geophys. Res. Lett.* (in press).

Solar Wind Isotopic Abundance Ratios of Ne, Mg, and Si Measured by SOHO/CELIAS/MTOF as a Diagnostic Tool for the Inner Corona, R. Kallenbach, F.M. Ipavich, H. Kucharek, P. Bochsler, A. B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, H. Grünwaldt, M. Hilchenbach, and D. Hovestadt. Manuscript submitted to *Physics and Chemistry of the Earth*, in press, 1999.

On the Solar Wind Composition during the November 1997 Solar Particle Events: WIND/MASS Observations, R.F. Wimmer-Schweingruber, O. Kern, and D.C. Hamilton., *Geophys. Res. Lett.*, 26, No. 23, 3541, 1999.

Composition and elemental abundance variations in the solar atmosphere and solar wind., Mason H., and Bochsler P., In: *Coronal holes and solar wind acceleration (Proceedings of the SOHO 7 Workshop, Northeast Harbor*, J. L. Kohl and S. R. Cranmer eds.) Kluwer Academic Publishers, 105, 1999.

Direct evidence of the interstellar gas flow velocity in the pickup ion cut-off as observed with SOHO CELIAS CTOF, E. Möbius, Y. Litvinenko, H. Grünwaldt, M.R. Aellig, A. Bogdanov, F.M. Ipavich, P. Bochsler, M. Hilchenbach, D. Judge, B. Klecker, M.A. Lee, and H. Ogawa, *Geophysical Research Letters*, 26, No. 20, 3181, 1999.

Determination of the Argon Isotopic Ratio of the Solar Wind using SOHO/CELIAS/MTOF, J.M. Weygand, F.M. Ipavich, P. Wurz, J.A. Paquette, and P. Bochsler, *Proceedings of 8th SOHO Workshop*, ESA SP-446, 1999.

New Observations of Long-Term Solar Soft X-Ray and EUV Fluxes from SOHO CELIAS/SEM Instrument, P. Gangopadhyay, H.S. Ogawa, D.R. McMullin, D.L. Judge, F. Ipavich, A.B. Galvin and M. Hilchenbach, *J. Geophys. Res..*, 28321, 1999.

Absolute Solar 30.4 nm Flux From Sounding Rocket Observations During The Solar Cycle 23 Minimum, D.L. Judge, D.R. McMullin, and H.S. Ogawa, *J. Geophys. Res.h.*, 1999.

Magnesium Isotopic Abundance in Slow and Coronal Hole Associated Solar Wind: SOHO/CELIAS/MTOF Measurements. H. Kucharek, F.M. Ipavich, R. Kallenbach, B. Klecker, H. Grünwaldt, M. R. Aellig, and P. Bochsler, *Proceedings of 8th SOHO Workshop*, ESA SP-446, 1999.

Observation of Suprathermal Helium at 1 AU: Charge States and Abundances in CIRs, M. Hilchenbach, H. Gruenwaldt, H. Kucharek, B. Klecker, F. M. Ipavich, A. B. Galvin, and R. Kallenbach, *Solar Wind 9, AIP conference proceedings 471*, New York, Editors: S. R. Habbal, R. Esser, J. Hollweg, and P. Isenberg, 1999.

The solar wind ionic charge states during the Ulysses pole-to-pole pass, Y.-K. Ko, G. Gloeckler, C. M. S. Cohen, and A. B. Galvin, *Solar Wind 9, AIP conference proceedings* 471, New York, Editors: S. R. Habbal, R. Esser, J. Hollweg, and P. Isenberg, 1999.

The Fe/O Elemental Abundance Ratio in the Solar Wind, M.R. Aellig, H. Holweger, P. Bochsler, P. Wurz, H. Grünwaldt, S. Hefti, H. Holweger, and B. Klecker, *Solar Wind 9, AIP conference proceedings 471*, New York, Editors: S. R. Habbal, R. Esser, J. Hollweg, and P. Isenberg, 255, 1999.

The Influence of Superathermal Electrons on the Derivation of Coronal Electron Temperatures from Solar Wind Motor Ion Charge States, M.R. Aellig, P. Bochsler, H. Grünwaldt, S. Hefti, P. Wurz, M. Hilchenbach, D. Hovestadt, F.M. Ipavich, and F. Gliem, *Phys. Chem. Earth (C)*, 24, No. 4, 407, 1999.

O5+ Observation From Various Solar Wind Speeds, H. Grünwaldt, M. Hilchenbach, M. R. Aellig, F. M. Ipavich, K. -U. Reiche, and B. Wilken, *Solar Wind 9, AIP conference proceedings 471*, New York, Editors: S. R. Habbal, R. Esser, J. Hollweg, and P. Isenberg, 259-262, 1999.

The Fe/O Elemental Abundance Ratio in the Solar Wind as observed with SOHO/CELIAS/CTOF, M.R. Aellig, S.Hefti, H.Grünwaldt, P.Bochsler, P.Wurz, and F.M.Ipavich, D.Hovestadt, *J. Geophys. Res.*, 104, 24, 1999.

Isotopes in the Solar Wind: New Results from ACE, SOHO, and WIND Robert Wimmer-Schweingruber, P. Bochsler, and P. Wurz, *Solar Wind 9, AIP conference proceedings 471*, New York, Editors: S. R. Habbal, R. Esser, J. Hollweg, and P. Isenberg, 147, 1999.

The Iron, Silicon, and Oxygen Abundance in the Solar Wind Measured with SOHO/CELIAS/MTOF, P. Wurz, M.R. Aellig, F.M. Ipavich, S. Hefti, P. Bochsler, A.B. Galvin, H. Grüenwaldt, M. Hilchenbach, F. Gliem, and D. Hovestadt. *Physics and Chemistry of the Earth*, 24, 421, 1999.

Kinetic Properties of Solar Wind Minor Ions and Protons Measured with SOHO/CELIAS, Hefti, H. Grünwaldt, F.M. Ipavich, P. Bochsler, D. Hovestadt, M.R. Aellig, M. Hilchenbach, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, B. Klecker, E. Marsch, E. Möbius, M. Neugebauer, and P. Wurz., *J. Geophys. Res.* 103, 29697, 1998.

Isotopic Composition of Solar Wind Nitrogen: First In-Situ Determination by CELIAS/MTOF on board SOHO, R. Kallenbach, J. Geiss, F. M. Ipavich, G. Gloeckler, P. Bochsler, F. Gliem, M. Hilchenbach, and D. Hovestadt, *ApJ*, 507, L185, 1998.

Magnesium Isotopic Composition as Observed with the CELIAS/MTOF Experiment on the SOHO Spacecraft, H. Kucharek, F. M. Ipavich, R. Kallenbach, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hillchenbach, W. I. Axford, H. Balsiger, A. Bürgi, M. A. Coplan, A. B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, K. C. Hsieh, D. J. Judge, B. Klecker, M. A. Lee, S. Livi, G. G. Managaze, E. Marsch, E. Möbius, M. Neugebauer, H. S. Ogawa, K. -U. Reiche, M. Scholer, M. I. Verigin, B. Wilken, and P. Wurz, *Journal of Geophysical Research*, 103, 26805, 1998.

Observations of Galactic Cosmic Ray Modulation during Earth-Directed Coronal Mass Ejections, P. Mäkelä, J. Torsti, M. Teittinen, E. Valtonen, E. Riihonen, and F.M. Ipavich, *Geophys. Res. Lett.* 25, 2951, 1998.

The Solar Minimum Active Region 7978, its X2.6/1B Flare, CME, and Interplanetary Shock Propagation of 9 July 1996, M. Dryer, M.D. Andrews, H. Aurass, C. Deforest, A. B. Galvin, H. Garcia, F.M. Ipavich, M. Karlicky, A. Kiplinger, A. Klassen, R. Meisner, S.E. Paswaters, Z. Smith, S.J. Tappin, B.J. Thompson, S.I. Watari, D.J. Michels, G.E. Brueckner, R.A. Howard, M.J. Koomen, P. Lamy, G. Mann, K. Arzner, and R. Schwenn, *Solar Physics 181*, 151-183, 1998.

A Crossroads for European Solar & Heliospheric Physics P. Bochsler, *Scientific achievements of SOHO: In situ solar wind*, Tenerife, March 23-27 1998. ESA SP-417, 35, 1998.

Fractionation of Si, Ne, and Mg isotopes in the solar wind as measured by SOHO/CELIAS/MTOF. R. Kallenbach, F. M. Ipavich, H. Kucharek, P. Bochsler, A. B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, H. Grünwaldt, S. Hefti, D. Hovestadt, and M. Hilchenbach, *Space Sci. Rev.*, 85, 357, 1998.

Detection of 55-80 keV Hydrogen Atoms of Heliospheric Origin by CELIAS/HSTOF on SOHO, M. Hilchenbach, K.C. Hsieh, D. Hovestadt, B. Klecker, H. Grüwaldt, P. Bochsler, F.M. Ipavich, A. Bürgi, E. Möbius, F. Gliem, W.I. Axford, H. Balsiger, W. Bornemann, M.A. Coplan, A.B. Galvin, J. Geiss, G. Gloeckler, S. Hefti, D.L. Judge, R. Kallenbach, P. Laeverenz, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, M. Neugebauer, H.S. Ogawa, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, *Astrophys. J.* 503, 916, 1998.

Kinetic Properties of Solar Wind Minor Ions and Protons Measured with SOHO/CELIAS, S. Hefti, H. Grünwaldt, F. M. Ipavich, P.Bochsler, D. Hovestadt, M. R. Aellig, M. Hilchenbach, A. B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, B. Klecker, E. Marsch, E. Möbius, M. Neugebauer, and P. Wurz, *J. Geophy. Res.*, 103, 29697, 1998.

Iron Freeze-In Temperatures Measured by SOHO/CELIAS/CTOF. M.R. Aellig, H. Grünwaldt, P. Bochsler, P. Wurz, S. Hefti, R. Kallenbach, F.M. Ipavich, W. I. Axford, A. Balsiger, A. Bürgi, M. A. Coplan, A. B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, M. Hilchenbach, D. Hovestadt, K. C. Hsieh, B. Klecker, M. A. Lee, S. Livi, G. G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, K.-U. Reiche, M. Scholer, M. I. Verigin, and B. Wilken, *J. Geophys. Res.* 103, 17215, 1998.

Solar Wind Measurements with SOHO: The CELIAS/MTOF Proton Monitor. F.M. Ipavich, A.B. Galvin, S.E. Lasley, J.A. Paquette, S. Hefti, K.-U. Reiche, M.A. Coplan,

Magnesium Isotopes in the Solar Wind as Observed with the MTOF Sensor on the CELIAS Experiment Onboard the SOHO Spacecraft, H. Kucharek, F.M. Ipavich, R. Kallenbach, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, W.I. Axford, H. Balsiger, A. Bürgi, M.A. Coplan, A.B. Galvin, J. Geiss, G. Gloeckler, K.C. Hsieh, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, P. Wurz, *Proceedings of the 5th SOHO Workshop*, ESA SP-404, 473, 1997.

The Corona and Solar Wind near Minimum Activity, P. Bochsler, Particles in the solar wind. *Proceedings of the Fifth SOHO Workshop*, ESA SP-404, 113, 1997.

The Sun at Minimum Activity: Results from the CELIAS Experiment on SOHO, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F.M. Ipavich, M.R. Aellig, W.I. Axford, H. Balsiger, A. Bogdanov, A. Bürgi, M.A. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, S. Hefti, K.C. Hsieh, D.L. Judge, R. Kallenbach, B. Klecker, H. Kucharek, S.E. Lasley, M.A. Lee, Yu. Litvinenko, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, H.S. Ogawa, J.A. Paquette, K.-U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, P. Wurz, *Proceedings of the 5th SOHO Workshop*, ESA SP-404, 37, 1997.

Solar Wind Iron Charge States Observed With High Time Resolution with SOHO/CELIAS/CTOF, M.R. Aellig, H. Grünwaldt, P. Bochsler, P. Wurz, S. Hefti, R. Kallenbach, F.M. Ipavich, D. Hovestadt, M. Hilchenbach, and the CELIAS Team, *Proceedings of the 5th SOHO Workshop*, ESA SP-404, 157, 1997.

Isotopic Composition of Solar Wind Neon Measured by CELIAS/MTOF onboard SOHO, R. Kallenbach, F.M. Ipavich, P. Bochsler, S. Hefti, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, W.I. Axford, H. Balsiger, A. Bürgi, M.A. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, K.C. Hsieh, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, K.U. Reiche, M. Scholer, M.I. Verigin, B. Wilken, P. Wurz, , *J. Geophys. Res.*, 102, 26895, 1997.

Venus Tail Ray Observation near Earth, H. Grünwaldt, M. Neugebauer, M. Hilchenbach, P. Bochsler, D. Hovestadt, A. Bürgi, F. M. Ipavich, K.-U. Reiche, W. I. Axford, H. Balsiger, A. B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, K. C. Hsieh, R. Kallenbach, B. Klecker, S. Livi, M. A. Lee, G. G. Managadze, E. Marsch, E. Möbius, M. Scholer, M. I. Verigin, B. Wilken, P. Wurz, *Geophys. Res. Lett.*, 24, 2557, 1997.

First Solar EUV Irradiances Obtained from SOHO by the CELIAS/SEM, D.L. Judge, D.R. McMullin, H.S. Ogawa, D. Hovestadt, B. Klecker, M. Hilchenbach, E. Möbius, L.R. Canfield, R.E. Vest, R. Watts, C. Tarrio, M. Kühne, and P. Wurz, *Solar Physics*, 177, 161, 1997.

Isotope Fractionation in Solar Energetic Particles by Stochastic Acceleration, R. Kallenbach, ESA SP-373, 191, 1994.

Fractionation of Nitrogen Isotopes in Solar Energetic Particles, P. Bochsler and R. Kallenbach, *Meteoritics*, 29, 653, 1994.

G. Gloeckler, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F. Gliem, W. I. Axford, A. Balsiger, A. Bürgi, J. Geiss, K. C. Hsieh, R. Kallenbach, B. Klecker, M. A. Lee, G. G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, M. Scholer, M. I. Verigin, B. Wilken, P. Wurz, *J. Geophys. Res.*, 103, 17205, 1998.

Elemental Composition of the January 6, 1997, CME, P. Wurz, F.M. Ipavich, A.B. Galvin, P. Bochsler, M.R. Aellig, R. Kallenbach, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, W.I. Axford, H. Balsiger, A. Bürgi, M.A. Coplan, J. Geiss, F. Gliem, G. Gloeckler, S. Hefti, K.C. Hsieh, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, K.U. Reiche, M. Scholer, M.I. Verigin, and B. Wilken, *Geophys. Res. Lett.* 25(14), 2557, 1998.

The Solar Minimum Active Region 7978, its X2.6/1B Flare, CME, and Interplanetary Shock Propagation of 9 July 1996, M. Dryer, M. D. Andrews, H. Aurass, C. DeForest, A. B. Galvin, H. Garcia, F. M. Ipavich, M. Karlicky, A. Kiplinger, A. Klassen, R. Meisner, S. E. Paswaters, Z. Smith, S. J. Tappin, B. J. Thompson, S. I. Watari, D. J. Michels, G.E. Brueckner, R. A. Howard, M. J. Koomen, P. Lamy, G. Mann, K. Arzner, and R. Schwenn, *Solar Physics* 181, 159, 1998.

Solar Wind Minor Ion Charge States Observed With High Time Resolution With SOHO/CELIAS/CTOF, M.R. Aellig, H. Grünwaldt, P. Bochsler, S. Hefti, P. Wurz, R. Kallenbach, F.M. Ipavich, D. Hovestadt, M. Hilchenbach, and the CELIAS Team, *Proceedings of the 31st ESLAB Symposium*, ESA-SP 415, 27, 1998.

Elemental Composition Before, during, and After the January 6, 1997, CME Event Measured by CELIAS/SOHO, P. Wurz, F.M. Ipavich, A.B. Galvin, P. Bochsler, M.R. Aellig, R. Kallenbach, D. Hovestadt, A. Grünwaldt, M. Hilchenbach, W.I. Axford, H. Balsiger, A. Bürgi, M.A. Coplan, J. Geiss, F. Gliem, G. Gloeckler, S. Hefti, K.C. Hsieh, B. Klecker, M.A. Lee, S. Livi, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, K.U. Reiche, M. Scholer, M.I. Verigin, and B. Wilken, *Proceedings of the 31st ESLAB Symposium*, ESA-SP 415, 395, 1998.

Limits on the Fractionation of Isotopes in the Solar Wind as Observed with SOHO/CELIAS/MTOF, R. Kallenbach, P. Bochsler, F.M. Ipavich, A.B. Galvin, R. Bodmer, S. Hefti, H. Kucharek, F. Gliem, H. Grünwaldt, M. Hilchenbach, B. Klecker, D. Hovestadt, and the CELIAS Team, *Proceedings of the 31st ESLAB Symposium*, ESA-SP 415, 33, 1998.

First year continuous solar EUV Irradiances from SOHO by the CELIAS/SEM during 1996 solar minimum, H.S. Ogawa, D.L. Judge, D.R. McMullin, P. Gangopadhyay, and A.B. Galvin, *J. Geophys. Res.*, 103, 1, 1998.

Solar Wind Measurements With SoHO: the CELIAS/MTOF Proton Monitor, F.M. Ipavich, A.B. Galvin, S.E. Lasley, J.A. Paquette, S. Hefti, K.-U. Reiche, M.A. Coplan, G. Gloeckler, P. Bochsler, D. Hovestadt, H. Grünwaldt, M. Hilchenbach, F. Gliem, W.I. Axford, H. Balsiger, A. Bürgi, J. Geiss, K.C. Hsieh, R. Kallenbach, B. Klecker, M.A. Lee, G.G. Managadze, E. Marsch, E. Möbius, M. Neugebauer, M. Scholer, M.I. Verigin, B. Wilken, and P. Wurz, J. Geophys. Res., 103, 17205, 1998.

Isotopic Composition of Solar Wind Calcium: First In Situ Measurement by CELIAS/MTOF on board SOHO, R. Kallenbach, F.M. Ipavich, P. Bochsler, S. Hefti, P. Wurz, M.R. Aellig, M.A. Coplan, A.B. Galvin, J. Geiss, F. Gliem, G. Gloeckler, H. Grünwaldt, M. Hilchenbach, D. Hovestadt, and B. Klecker, *ApJ Letters*, 498, L75, 1997.

# Related Presentations

Oxygen 16 to Oxygen 18 abundance ratio in the solar wind observed by Wind/MASS, M. R. Collier, D. C. Hamilton, G. Gloeckler, G. Ho, P. Bochsler, R. Bodmer, and R. Sheldon, *J. Geophys. Res.* 103, 7, 1998.

Particles in the Solar Wind, P. Bochsler, 5th SOHO Workshop, 17-20 June, 1997, Oslo, Norway.